Index to Volume 77 **Author Index**

Page numbers of errata are in italic.

Abecassis, J. See E. Charun, 265

See R. Pujol, 421

Adachi, T., K. Watanabe, and T. Mitsunaga. Characterization of thiaminbinding protein from wheat germ, 578

Åman, P. See A. A. M. Andersson, 463

Ammar, K., W. E. Kronstad, and C. F. Morris. Breadmaking quality of selected durum wheat genotypes and its relationship with high molecular weight glutenin subunits allelic variation and gluten protein polymeric composition,

Anderson, A. K., and P. K. W. Ng. Changes in disulfide and sulfhydryl contents and electrophoretic patterns of extruded wheat flour proteins, 354 Andersson, A. A. M., R. Andersson, and P. Aman. Air classification of barley flours, 463

Andersson, R. See A. A. M. Andersson, 463

Ando, H. See H. Tang, 27

Andrews, L. C. See C. S. Gaines, 187

Arendt, E. K. See C. M. O'Brien, 111

See P. Crowley, 370

Artz, W. E. See S. M. Mahungu, 220 Autran, J. C. See M.-H. Morel, 685

Åvall, A.-K. See E. Bertoft, 657

Baik, B.-K. See G. G. Mikhaylenko, 507

Baik, M.-Y., and P. Chinachoti. Moisture redistribution and phase transitions during bread staling, 484

Bailey, T. B. See A. E. McPherson, 320

Bar-L'Helgouac'h, C. See M.-H. Morel, 685

Barragán Delgado, M. L., and S. O. Serna Saldivar. Production and nutritional evaluation of liquefied weaning foods from malted sorghum, quality protein maize, and other cereals, 652

Barrett, A. H., A. V. Cardello, L. Mair, P. Maguire, L. L. Lesher, M. Richardson, J. Briggs, and I. A. Taub. Textural optimization of shelf-stable bread: Effects of glycerol content and dough-forming technique, 169

M. Tsoubeli, P. Maguire, Tan, N. B., K. Conca, Y. Wang, B. Porter, and I. Taub. Textural stability of intermediate-moisture extrudates: Effects of formulation, 784

Barton, F. E., II, D. S. Himmelsbach, A. M. McClung, and E. T. Champagne. Rice quality by spectroscopic analysis: Precision of three spectral regions, 669 Batey, I. L., and B. M. Curtin. Effects on pasting viscosity of starch and flour from different operating conditions for the Rapid Visco Analyser, 754

Bechtel, D. B., and J. Wilson. Variability in a starch isolation method and automated digital image analysis system used for the study of starch size distributions in wheat flour, 401

—. See C. S. Gaines, 163 Beers, K. W. See B. J. Lloyd, 551

Bejosano, F. P. See M. Cepeda, 489

Bekes, F. See O. R. Larroque, 451

See S. Uthayakumaran, 731, 737, 744

See W. S. Veraverbeke, 582, 589 Bennink, M. R. See V. E. A. Rinaldi, 237

Bergman, C. J., J. T. Delgado, R. Bryant, C. Grimm, K. R. Cadwallader, and B. D. Webb. Rapid gas chromatographic technique for quantifying 2-acetyl-1- pyrroline and hexanal in rice (Oryza sativa, L.), 454

Bertoft, E., C. Boyer, R. Manelius, and A.-K. Åvall. Observations on the αamylolysis pattern of some waxy maize starches from inbred line Ia453, 657

See R. Manelius, 345 Bett, K. L. See J.-F. Meullenet, 512

Bettge, A. D., M. J. Giroux, and C. F. Morris. Susceptibility of waxy starch granules to mechanical damage, 750

, and C. F. Morris. Relationships among grain hardness, pentosan fractions, and end-use quality of wheat, 241

Bhatty, R. S. See G. H. Zheng, 140 Bietz, J. A. See J. L. Robutti, 91

Black, C. K., J. F. Panozzo, C. L. Wright, and P. C. Lim. Survey of white salted noodle quality characteristics in wheat landraces, 468

Blanshard, J. M. V. See I. A. Farhat, 202

Borrás, F. S. See J. L. Robutti, 24, 91

Boyer, C. See E. Bertoft, 657

Boyer, J. E., Jr. See J. Suroso, 808

Briggs, J. See A. H. Barrett, 169

Bryant, R. See C. J. Bergman, 454

Budi Santosa, F. X., and G. W. Padua. Thermal behavior of zein sheets plasticized with oleic acid, 459

Buescher, R. W. See A. L. Dillahunty, 541

Buriak, P. See J. Wahjudi, 640

-. See P. Yang, 44, 128

Cadwallader, K. R. See C. J. Bergman, 454

Campbell, M. R., J. Sykes, and D. V. Glover. Classification of single- and doublemutant corn endosperm genotypes by near-infrared transmittance spectroscopy,

Cao, T. K. See G. H. Robertson, 439

Capocchi, A., L. Galleschi, and F. Saviozzi. Note: Isolation of wheat high molecular weight glutenin subunits from durum wheat, 105

Carbone, L. See E. A. Tosi, 699

Cardello, A. V. See A. H. Barrett, 169

Cepeda, M., R. D. Waniska, L. W. Rooney, and F. P. Bejosano. Effects of leavening acids and dough temperatures in wheat flour tortillas, 489

Cerletti, P. See D. Scardone, 602

Champagne, E. T. See F. E. Barton II, 669

See B. G. Lyon, 64

See J.-F. Meullenet, 512 Chan, R. See F. M. DuPont, 607

Charun, E., J. Abecassis, A.-S. Contamine, T.-M. Roulland, B. Vergnes, and M.-H. Morel. Effects of temperature and mechanical input on semisweet biscuit (cookie) quality and dough characteristics, 265

Chatakanonda, P., S. Varavinit, and P. Chinachoti. Relationship of gelatinization and recrystallization of cross-linked rice to glass transition temperature, 315

Chaurand, M. See R. Pujol, 421

Cheryan, M. See R. Shukla, 724

Chien, J. T. See Y .- C. Lai, 544

Chinachoti, P. See M.-Y. Baik, 478

See P. Chatakanonda, 315 Cho, K.-Y. See S.-Y. Won, 309

Choi, W. S. See S.-Y. Won, 309

Chow, F. I. See T. S. Kahlon, 518, 673

Chun, J. K. See K. M. Chung, 567

Chung, D. S. See F. C. Wang, 478

Chung, K. M., T. W. Moon, and J. K. Chun. Influence of annealing on gel properties of mung bean starch, 567

Chung, O. K. See J. B. Ohm, 556

. See A. Sayaslan, 248

Conca, K. See A. Barrett, 784

Conde-Petit, B. See C. Zweifel, 645 Contamine, A.-S. See E. Charun, 265

Courtin, C. M. See U. Elofsson, 679

Crowley, P., H. Grau, and E. K. Arendt. Influence of additives and mixing time on crumb grain characteristics of wheat bread, 370

Cubadda, R. See E. Marconi, 133

Cuniberti, M. See E. A. Tosi, 699

Curtin, B. M. See I. L. Batey, 754 Czuchajowska, Z. See G. G. Mikhaylenko, 507

Dailey, O. D., Jr. Note: Variability in water absorption of germ and endosperm during laboratory steeping of a yellow corn hybrid, 721

Damerval, C. See J. Landry, 620

Daniels, M. J. See J.-F. Meullenet, 259

Dehlon, P. See M.-H. Morel, 685

Delcour, J. A. See U. Elofsson, 679

See W. S. Veraverbeke, 582, 589 Delgado, J. T. See C. J. Bergman, 454

Delhaye, S. See J. Landry, 620

Del Nobile, M. A., and M. Massera. Modeling of water sorption kinetics in spaghetti during overcooking, 615

Delwiche, S. R., and W. R. Hruschka. Note: Protein content of bulk wheat from near-infrared reflectance of individual kernels, 86

Denery-Papini, S. See C. Larré, 121

Descamps, M. See I. A. Farhat, 202

Deshayes, G. See C. Larré, 121 See J. Lefebvre, 193 Desserme, C. See C. Larré, 121

DeVor, R. E. See R. Shukla, 724 Dexter, J. E. See B. C. Morgan, 286

Dillahunty, A. L., T. J. Siebenmorgen, R. W. Buescher, D. E. Smith, and A. Mauromoustakos. Effect of moisture content and temperature on respiration rate of rice, 541

Doehlert, D. C., and M. S. McMullen. Genotypic and environmental effects on oat milling characteristics and groat hardness, 148

Donelson, J. R., C. S. Gaines, and P. L. Finney. Baking formula innovation to eliminate chlorine treatment of cake flour, 53

Donelson, T. See C. S. Gaines, 163 Doner, L. W. See V. Singh, 560

Dowell, F. E. Differentiating vitreous and nonvitreous durum wheat kernels by using near-infrared spectroscopy, 155 Drozdek, K. A. See S. M. Mahungu, 220

Du, L. See P. Yang, 128

DuPont, F. M., W. H. Vensel, R. Chan, and D. D. Kasarda. Characterization of the 1B-type ω-gliadins from Triticum aestivum cultivar Butte, 607

Eckhoff, S. R. See S. Mehra, 209

See V. Singh, 560, 665, 692

See J. Wahjudi, 640 See D. Wang, 525

See P. Yang, 44, 128, 529

Ekholm, P., L. Virkki, M. Ylinen, L. Johansson, and P. Varo. Effects of natural chelating agents on the solubility of some physiologically important mineral elements in oat bran and oat flakes, 562

Eliasson, A.-C. See U. Elofsson, 679

See H. Larsson, 633

Elofsson, U., A.-C. Eliasson, M. Wahlgren, A.-M. A. Loosveld, C. M. Courtin, and J. A. Delcour. Adsorption studies of interaction between water-extractable nonstarch polysaccharides and prolamins in cereals, 679

Engle, D. A. See C. F. Morris, 77 Escher, F. See C. Zweifel, 645

Faller, J. F. See S. M. Mahungu, 220

Fang, Q., and M. A. Hanna. Functional properties of polylactic acid starchbased loose-fill packaging foams, 779

Farhat, I. A., J. M. V. Blanshard, M. Descamps, and J. R. Mitchell. Effect of sugars on retrogradation of waxy maize starch-sugar extrudates, 202

Favier, J. F. See E. L. Lazaro, 717 Ferrer, M. E. See J. L. Robutti, 24, 91

Fessas, D. See M. Riva, 433

Finney, P. L. See J. R. Donelson, 53 . See C. S. Gaines, 163, 187

Fitzgerald, M. A. See A. J. Lisle, 627

Flores, R. A. See J. Suroso, 808

Forlani, F. See D. Scardone, 602

Forssell, P. See M. Lauro, 595

Fraignier, M.-P., N. Michaux-Ferrière, and K. Kobrehel. Distribution of peroxidases in durum wheat (Triticum durum), 11

Gaines, C. S., P. L. Finney, and L. C. Andrews. Developing agreement between very short flow and longer flow test wheat mills, 187

M. Ö. Raeker, M. Tilley, P. L. Finney, J. D. Wilson, D. B. Bechtel, R. J. Martin, P. A. Seib, G. L. Lookhart, and T. Donelson. Associations of starch gel hardness, granule size, waxy allelic expression, thermal pasting, milling quality, and kernel texture of 12 soft wheat cultivars, 163

See J. R. Donelson, 53

Galleschi, L. See A. Capocchi, 105 Gaosong, J., and T. Vasanthan. Effect of extrusion cooking on the primary structure and water solubility of \beta-glucans from regular and waxy barley,

Gelroth, J. A. See G. S. Ranhotra, 159, 293

Gianibelli, M. C. See O. R. Larroque, 448

Gilbert, S. M. See A. D. L. Humphris, 107

Giroux, M. J. See A. D. Bettge, 750

Glenn, G. M. See W. J. Orts, 18

Glover, D. V. See M. R. Campbell, 774

Gomez Sanchez, M. See O. R. Larroque, 448

Gras, P. W. See S. Uthayakumaran, 731, 737

Grau, H. See P. Crowley, 370

See C. M. O'Brien, 111

Graybosch, R. A., G. Guo, and D. R. Shelton. Note: Aberrant falling numbers of waxy wheats independent of α-amylase activity, 1

Graziano, M. See E. Marconi, 133

Griffin, V. K. See J.-F. Meullenet, 259

Grimm, C. See C. J. Bergman, 454 Guan, F. See P. A. Seib, 816 Guo, G. See R. A. Gravbosch, 1

Haken, A. E. See V. Singh, 665, 692

See P. Yang, 44

Hamaker, B. R. See C. P. Huang, 343

Han, X. Z. See C. P. Huang, 343 Hankins, J.-A. See J.-F. Meullenet, 259

Hanna, M. A. See Q. Fang, 779

Harada, O., E. D. Lysenko, and K. R. Preston. Effects of commercial hydrolytic enzyme additives on Canadian short process bread properties and processing characteristics, 70

Hare, R. A. See M. J. Sissons, 4

Hatcher, D. W., and S. J. Symons. Assessment of oriental noodle appearance as a function of flour refinement and noodle type by image analysis, 181

-. Image analysis of Asian noodle appearance: Impact of hexaploid wheat with a red seed coat, 388

-. Influence of sprout damage on oriental noodle appearance as and assessed by image analysis, 380

Hejlsoe-Kohsel, E. See C. P. Huang, 343

Hicks, K. B. See V. Singh, 560, 665, 692

Himmelsbach, D. S. See F. E. Barton II, 669

Hizukuni, S. See M. Tako, 473

Hosomi, K. See M. Seguchi, 339

Hruschka, W. R. See S. R. Delwiche, 86

Huang, C. P., E. Hejlsoe-Kohsel, X. Z. Han, and B. R. Hamaker. Note: Proteolytic activity in sorghum flour and its interference in protein analysis, 343

Hugo, L. F., L. W. Rooney, and J. R. N. Taylor. Malted sorghum as a func-

tional ingredient in composite bread, 428 Humphris, A. D. L., T. J. McMaster, M. J. Miles, S. M. Gilbert, P. R. Shewry, and A. S. Tatham. Atomic force microscopy (AFM) study of interactions of HMW subunits of wheat glutenin, 107

Imai, T. See M. Seguchi, 339 Ingelin, M. E. See J. Suchy, 39

Jackson, D. S. See M. G. Osman, 101

Jackson, R. See M. J. Sissons, 4

Jane, J.-L. See K. Koch, 115

See A. E. McPherson, 320, 326

Jeffers, H. C. See C. F. Morris, 77

Johansson, E. See H. Larsson, 633 Johansson, L. See P. Ekholm, 562

Johnston, D. B. See V. Singh, 560

Kaarlehto, T., and H. Salovaara. Effect of kiln drying on falling number of oats, 177

Kahlon, T. S., and F. I. Chow. In vitro binding of bile acids by rice bran, oat bran, wheat bran, and corn bran, 518

-. Lipidemic response of hamsters to rice bran, uncooked or processed white and brown rice, and processed corn starch, 673

Kasarda, D. D. See F. M. DuPont, 607

Kauffmann, D. See J.-F. Meullenet, 512

Kavale, S. See A. R. Wooding, 791, 798

Ke, T., and X. Sun. Physical properties of poly(lactic acid) and starch composites with various blending ratios, 761 Keentok, M. See S. Uthayakumaran, 744

Keogh, M. K. See C. M. O'Brien, 111

Kerr, W. L., C. D. W. Ward, K. H. McWatters, and A. V. A. Resurreccion. Effect of milling and particle size on functionality and physicochemical properties of cowpea flour, 213

Kidwell, K. K. See G. G. Mikhaylenko, 507

Kim, D. C. See S. S. Kim, 376

Kim, O. W. See S. S. Kim, 376

Kim, S. S., S. E. Lee, O. W. Kim, and D. C. Kim. Physicochemical characteristics of chalky kernels and their effects on sensory quality of cooked

Kim, Y. S. See F. C. Wang, 478

Knutson, C. A. See J. Robutti, 24

Kobrehel, K. See M.-P. Fraignier, 11

Koch, K., and J.-L. Jane. Morphological changes of granules of different starches by surface gelatinization with calcium chloride, 115

Kronstad, W. E. See K. Ammar, 230

Kunetz, C. F. See E. L. Suhendro, 96

Labat, E., M. H. Morel, and X. Rouau. Effects of laccase and ferulic acid on wheat flour doughs, 823

Lai, K. P., J. F. Steffe, and P. K. W. Ng. Average shear rates in Rapid Visco

Analyser (RVA) mixing system, 714

Lai, V. M.-F., S. Lu, and C. Lii. Molecular characteristics influencing retro-

gradation kinetics of rice amylopectins, 272 Lai, Y.-C., P.-H. Sung, and J. T. Chien. Evaluation of compatibility of rice starch and pectins by glass transition and sub-Tg endosperms and the effect of compatibility on gel viscosity and water loss, 544

Landry, J., S. Delhaye, and C. Damerval. Improved method for isolating and quantitating α-amino nitrogen as nonprotein, true protein, salt-soluble proteins, zeins, and true glutelins in maize endosperm, 620

Larré, C., S. Denery-Papini, Y. Popineau, G. Deshayes, C. Desserme, and J. Lefebvre. Biochemical analysis and rheological properties of gluten modified by transglutaminase, 121

See E. Linarès, 414

Larroque, O. R., M. C. Gianibelli, M. Gomez Sanchez, and F. MacRitchie. Note: Procedure for obtaining stable protein extracts of cereal flour and whole meal for size-exclusion HPLC analysis, 448

and F. Bekes. Note: Rapid size-exclusion chromatography analysis of molecular size distribution for wheat endosperm protein, 451

-. See W. S. Veraverbeke, 582, 589

Larsson, H., A.-C. Eliasson, E. Johansson, and G. Svensson. Influence of added starch on mixing of dough made with three wheat flours differing in high molecular weight subunit composition: Rheological behavior, 633

Lauro, M., K. Poutanen, and P. Forssell. Effect of partial gelatinization and lipid addition on \alpha-amylolysis of barley starch granules, 595

Lavenant, L. See J. Lefebvre, 193

Lazaro, E. L., and J. F. Favier. Note: Alkali debranning of sorghum and millet, 717

Ledesma-Osuna, A. I. See P. I. Torres, 702

Lee, E. Y., K. I. Lim, J. Lim, and S.-T. Lim. Effect of gelatinization degree and moisture content of extruded corn starch pellets on morphology and physical properties of microwave-expanded products, 769

Lee, S. E. See S. S. Kim, 376

Lefebvre, J., Y. Popineau, G. Deshayes, and L. Lavenant. Temperatureinduced changes in the dynamic rheological behavior and size distribution of polymeric proteins for glutens from wheat near-isogenic lines differing in HMW glutenin subunit composition, 193

See C. Larré, 121

Leinen, S. D. See G. S. Ranhotra, 293 Lemeste, M. See E. Linarès, 414 Lempereur, I. See R. Pujol, 421 Lens, J.- P. See V. Tropini, 333 Lesher, L. L. See A. H. Barrett, 169 Létang, C. See R. Pujol, 421 Leygue, J. P. See M.-H. Morel, 685 Li, B. H. See P. Yang, 128 Liang, X. See P. A. Seib, 816 Liang, Y. T. See P. A. Seib, 816 Lii, C. See V. M.-F. Lai, 272

Lim, H. S. See S.-Y. Won, 309 Lim, J. See E. Y. Lee, 769

Lim, K. I. See E. Y. Lee, 769 Lim, P. C. See C. K. Black, 468

Lim, S.-T. See E. Y. Lee, 769

See S.-Y. Won, 309 See S. You, 303

Linarès, E., C. Larré, M. Lemeste, and Y. Popineau. Emulsifying and foaming properties of gluten hydrolysates with an increasing degree of hydrolysis: Role of soluble and insoluble fractions, 414

Lindsay, M. P., and J. H. Skerritt. Immunocytochemical localization of gluten proteins uncovers structural organization of glutenin macropolymer, 360

Lingnert, H. See A. Rutgersson, 407

Lisle, A. J., M. Martin, and M. A. Fitzgerald. Chalky and translucent rice grains differ in starch composition and structure and cooking properties, 627

Lloyd, B. J., T. J. Siebenmorgen, and K. W. Beers. Effects of commercial processing on antioxidants in rice bran, 551

Lookhart, G. L. See C. S. Gaines, 163

Loosveld, A.-M. A. See U. Elofsson, 679

Lu, S. See V. M.-F. Lai, 272 Lukow, O. M. See J. Suchy, 39

Lupton, J. R. See S. M. Maier, 297

Lyon, B. G., E. T. Champagne, B. T. Vinyard, and W. R. Windham. Sensory and instrumental relationships of texture of cooked rice from selected cultivars and postharvest handling practices, 64

Lysenko, E. D. See O. Harada, 70

Mabille, F. See R. Pujol, 421 MacRitchie, F. See O. R. Larroque, 448 See A. R. Wooding, 798 Maguire, P. See A. H. Barrett, 169, 784

Mahungu, S. M., K. A. Drozdek, W. E. Artz, and J. F. Faller. Residence time distribution and barrel fill in pet food twin-screw extrusion cooking, 220

Maier, S. M., N. D. Turner, and J. R. Lupton. Serum lipids in hypercholesterolemic men and women consuming oat bran and amaranth products, 297 Mair, L. See A. H. Barrett, 169

Manelius, R., K. Nurmi, and E. Bertoft. Enzymatic and acidic hydrolysis of cationized waxy maize starch granules, 345

See E. Bertoft, 657

Marconi, E., M. Graziano, and R. Cubadda. Composition and utilization of barley pearling by-products for making functional pastas rich in dietary fiber and β-glucans, 133

Marks, B. P. See J.-F. Meullenet, 259 Martin, M. See A. J. Lisle, 627

Martin, R. J. See C. S. Gaines, 163

Massera, M. See M. A. Del Nobile, 615

Matsuki, J. See T. Sasaki, 58

Mauromoustakos, A. See A. L. Dillahunty, 541

See A. Perdon, 708

McAloon, A. J. See J. Wahjudi, 640 McClung, A. M. See F. E. Barton II, 669

See J.-F. Meullenet, 512

McDonough, C. M. See E. L. Suhendro, 96

McMaster, T. J. See A. D. L. Humphris, 107

McMullen, M. S. See D. C. Doehlert, 148

McPherson, A. E., T. B. Bailey, and J. Jane. Extrusion of cross-linked hydroxypropylated corn starches. I. Pasting properties, 320

and J. Jane. Extrusion of cross-linked hydroxypropylated corn starches. II. Morphological and molecular characterization, 326

McWatters, K. H. See W. L. Kerr, 213

Medina-Rodriguez, C. See P. I. Torres, 702 Mehra, S., V. Singh, M. E. Tumbleson, and S. R. Eckhoff. Effect of mill plate setting and number of dynamic steeping stages for an intermittent milling and dynamic steeping (IMDS) process for corn, 209

Meullenet, J.-F., E. T. Champagne, K. L. Bett, A. M. McClung, and D. Kauffmann. Instrumental assessment of cooked rice texture characteristics: A method for breeders, 512

B. P. Marks, J.-A. Hankins, V. K. Griffin, and M. J. Daniels. Sensory quality of cooked long-grain rice as affected by rough rice moisture content, storage temperature, and storage duration, 259

See C. Sitakalin, 501 Michaux-Ferrière, N. See M.-P. Fraignier, 11

Mikhaylenko, G. G., Z. Czuchajowska, B.-K. Baik, and K. K. Kidwell. Environmental influences on flour composition, dough rheology, and baking quality of spring wheat, 507

Mikola, M., and B. L. Jones. Characterization of oat endoproteinases that hydrolyze oat globulins, 572

Miles, M. J. See A. D. L. Humphris, 107

Mitchell, J. R. See I. A. Farhat, 202

Mitsunaga, T. See T. Adachi, 578

. See H. Tang, 27

Moon, T. W. See K. M. Chung, 567

Moreau, R. A. See V. Singh, 665, 692

Morel, M.-H., P. Dehlon, J. C. Autran, J. P. Leygue, and C. Bar-L'Helgouac'h. Effects of temperature, sonication time, and power settings on size distribution and extractability of total wheat flour proteins as determined by size-exclusion high-performance liquid chromatography, 685

See E. Charun, 265

. See E. Labat, 823

Morgan, B. C., J. E. Dexter, and K. R. Preston. Relationship of kernel size to flour water absorption for Canada Western Red Spring wheat, 286

Morris, C. F., H. C. Jeffers, and D. A. Engle. Effect of processing, formula and measurement variables on alkaline noodle color-toward an optimized laboratory system, 77

See K. Ammar, 230

See A. D. Bettge, 241, 750

Mua, J. P. See D. Sahai, 254

Mulder, W. J. See V. Tropini, 333

Neville, D. P. See C. M. O'Brien, 111 Newberry, M. See S. Uthayakumaran, 744 Ng, P. K. W. See A. K. Anderson, 354

See K. P. Lai, 714 See V. E. A. Rinaldi, 237

Nishiba, Y., T. Sato, and I. Suda. Convenient method to determine free fatty acid of rice using thin-layer chromatography and flame-ionization detection

Niu, Y. X. See P. Yang, 44 Nobes, G. A. R. See W. J. Orts, 18 Nurmi, K. See R. Manelius, 345

O'Brien, C. M., H. Grau, D. P. Neville, M. K. Keogh, W. J. Reville, and E. K. Arendt. Effects of microencapsulated high-fat powders on the empirical and fundamental rheological properties of wheat flour doughs, 111

Ohm, J. B., and O. K. Chung. NIR transmittance estimation of free lipid content and its glycolipid and digalactosyldiglyceride contents using wheat flour lipid extracts, 556

Ojeda, C. A., M. P. Tolaba, and C. Suárez. Note: Modeling starch gelatinization kinetics of milled rice flour, 145

Okot-Kotber, B. M. See G. S. Ranhotra, 159

Orts, W. J., G. M. Glenn, G. A. R. Nobes, and D. F. Wood. Wheat starch effects on the textural characteristics of puffed brown rice cakes, 18

Osborne, B. G. See M. J. Sissons, 4

Osman, M. G., D. Sahai, and D. S. Jackson. Note: Oil absorption characteristics of a multigrain extrudate during frying: Effect of extrusion temperature and screw speed, 101

Padua, G. W. See F. X. Budi Santosa, 459

Panozzo, J. F. See C. K. Black, 468 Percibaldi, M. See J. Robutti, 24

Perdon, A., T. J. Siebenmorgen, and A. Mauromoustakos. Glassy state transition and rice drying: Development of a brown rice state diagram, 708

Popineau, Y. See C. Larré, 121

See J. Lefebvre, 193

----. See E. Linarès, 414 Porter, B. See A. Barrett, 784

Poutanen, K. See M. Lauro, 595

Preston, K. R. See O. Harada, 70 —. See B. C. Morgan, 286

Pujol, R., C. Létang, I. Lempereur, M. Chaurand, F. Mabille, and J. Abecassis. Description of a micromill with instrumentation for measuring grinding characteristics of wheat grain, 421

Raeker, M. Ö. See C. S. Gaines, 163

Ranhotra, G. S., J. A. Gelroth, and S. D. Leinen. Utilization of calcium in breads highly fortified with calcium as calcium carbonate or as dairy calcium, 293

—, and B. M. Okot-Kotber. Stability and dietary contribution of vitamin E added to bread, 159

Rausch, K. D. See J. Wahjudi, 640

---. See P. Yang, 128

Ré, E. D. See E. A. Tosi, 699

Reinikainen, P. See A. Rutgersson, 407

Resurreccion, A. V. A. See W. L. Kerr, 213

Reville, W. J. See C. M. O'Brien, 111

Richardson, M. See A. H. Barrett, 169

Rinaldi, V. E. A., P. K. W. Ng, and M. R. Bennink. Effects of extrusion on dietary fiber and isoflavone contents of wheat extrudates enriched with wet okara, 237

Riva, M., D. Fessas, and A. Schiraldi. Starch retrogradation in cooked pasta and rice, 433

Robertson, G. H., T. K. Cao, and D. F. Wood. Effect of morphology of mechanically developed wheat flour and water on starch from gluten separation using cold ethanol displacement, 439

Robutti, J. L., F. S. Borrás, M. E. Ferrer, and J. A. Bietz. Grouping and identification of Argentine maize races by principal component analysis of zein reversed-phase HPLC data, 91

_______, ______, M. Percibaldi, and C. A. Knutson. Evaluation of quality factors in Argentine maize races, 24

Rooney, L. W. See M. Cepeda, 489

____. See L. F. Hugo, 428

----. See E. L. Suhendro, 96

Rossnagel, B. G. See G. H. Zheng, 140

Rouau, X. See E. Labat, 823

Roulland, T.-M. See E. Charun, 265

Rowe, M. See D. Sahai, 254

Rutgersson, A., V.-M. Toukkuri, P. Reinikainen, and H. Lingnert. Influence of hydrothermal treatment on lipid oxidation in barley, 407

Sahai, D., I. Surjewan, J. P. Mua, M. O. Buendia, M. Rowe, and D. S. Jackson. Dry matter loss during nixtamalization of a white corn hybrid: Impact of processing parameters, 254

Salovaara, H. See T. Kaarlehto, 177

Sarker, R. See F. L. Stoddard, 445

Sasaki, T., T. Yasui, and J. Matsuki. Effect of amylose content on gelatinization, retrogradation, and pasting properties of starches from waxy and nonwaxy wheat and their F1 seeds, 58

Sato, T. See Y. Nishiba, 223

Saviozzi, F. See A. Capocchi, 105

Sayaslan, A., O. K. Chung, P. A. Seib, and L. M. Seitz. Volatile compounds

in five starches, 248

Scardone, D., F. Forlani, and P. Cerletti. Accessibility of amino groups in gluten proteins studied by a combination of chemical labeling and immunochemical detection, 602

Schiraldi, A. See M. Riva, 433

Seguchi, M., T. Yasui, K. Hosomi, and T. Imai. Study of internal structure of waxy wheat starch granules by KI/I₂ solution, 339

Seib, P. A., X. Liang, F. Guan, Y. T. Liang, and H. C. Yang. Comparison of Asian noodles from some hard white and hard red wheat flours, 816

----. See F. C. Wang, 478

--- See X. J. Xie, 392, 696

Seitz, L. M. See A. Sayaslan, 248

Serna Saldivar, S. O. See M. L. Barragán Delgado, 652

Shelton, D. R. See R. A. Graybosch, 1

Shukla, R., M. Cheryan, and R. E. DeVor. Solvent extraction of zein from dry-milled com, 724

Shunk, R. J. See P. Yang, 44

Siebenmorgen, T. J. See A. L. Dillahunty, 541

—. See B. J. Lloyd, 551

---. See A. Perdon, 708

Silvestre, F. See V. Tropini, 333

Singh, V., L. W. Doner, D. B. Johnston, K. B. Hicks, and S. R. Eckhoff. Note: Comparison of coarse and fine corn fiber for corn fiber gum yields and sugar profiles, 560

—, R. A. Moreau, A. E. Haken, S. R. Eckhoff, and K. B. Hicks. Hybrid variability and effect of growth location on corn fiber yields and corn fiber

oil composition, 692

----, ----, K. B. Hicks. and S. R. Eckhoff. Effect of various acids and sulfites in steep solution on yields and composition of corn fiber and corn fiber oil, 665

— See S. Mehra, 209

Sissons, M. J., B. G. Osborne, R. A. Hare, S. A. Sissons, and R. Jackson. Application of the single-kernel characterization system to durum wheat testing and quality prediction, 4

Sissons, S. A. See M. J. Sissons, 4

Sitakalin, C., and J.-F. C. Meullenet. Prediction of cooked rice texture using extrusion and compression tests in conjunction with spectral stress strain analysis, 501

Skerritt, J. H. See M. P. Lindsay, 360

Smith, D. E. See A. L. Dillahunty, 541

Steffe, J. F. See K. P. Lai, 714

Stoddard, F. L., and R. Sarker. Rapid Communication: Characterization of starch in *Aegilops* species, 445

---. See S. Uthayakumaran, 731, 737, 744

---. See A. R. Wooding, 791, 798

Suárez, C. See C. A. Ojeda, 145

Suchy, J., O. M. Lukow, and M. E. Ingelin. Dough microextensibility method using a 2-g mixograph and a texture analyzer, 39

Suda, I. See Y. Nishiba, 223

Suhendro, E. L., C. F. Kunetz, C. M. McDonough, L. W. Rooney, and R. D. Waniska. Cooking characteristics and quality of noodles from food sorghum, 96

Sun, X. S. See T. Ke, 761

---. See Z. K. Zhong, 495

Sung, P.-H. See Y.-C. Lai, 544

Surjewan, I. See D. Sahai, 254

Suroso, J., R. A. Flores, and J. E. Boyer, Jr. Scarification and degermination of sorghum for grits production: Effects of hybrid and conditioning, 808

Svensson, G. See H. Larsson, 633 Sykes, J. See M. R. Campbell, 774

Symons, S. J. See D. W. Hatcher, 181, 380, 388

Takeda, Y. See H. Tang, 27

....... See Y. Yoshimoto, 279

Takenouchi, T. See Y. Yoshimoto, 279

Tako, M., and S. Hizukuri. Retrogradation mechanism of rice starch, 473

Tan, N. B. See A. Barrett, 784

Tang, H., H. Ando, K. Watanabe, Y. Takeda, and T. Mitsunaga. Some physicochemical properties of small-, medium-, and large-granule starches in fractions of waxy barley grain, 27

Tashiro, J. See Y. Yoshimoto, 279

Tatham, A. S. See A. D. L. Humphris, 107

Taub, I. A. See A. H. Barrett, 169, 784 Taylor, J. R. N. See L. F. Hugo, 428

Tilley, M. See C. S. Gaines, 163

Tolaba, M. P. See C. A. Ojeda, 145

Torres, P. I., L. Vazquez-Moreno, A. I. Ledesma-Osuna, and C. Medina-Rodriguez. Contribution of hydrophobic soluble gluten proteins, fractionated by hydrophobic interaction chromatography in highly acetylated agarose, to dough rheological properties, 702

Tosi, E. A., E. D. Ré, L. Carbone, and M. Cuniberti. Breadmaking quality estimation by fast spectrophotometric method, 699

Toukkuri, V.-M. See A. Rutgersson, 407

Tropini, V., J.-P. Lens, W. J. Mulder, and F. Silvestre. Cross-linking of wheat gluten using a water-soluble carbodiimide, 333

Tsoubeli, M. See A. Barrett, 784

Tumbleson, M. E. See S. Mehra, 209

See J. Wahjudi, 640 -. See P. Yang, 44

Turner, N. D. See S. M. Maier, 297

Tyler, R. T. See G. H. Zheng, 140

Uthayakumaran, S., M. Newberry, M. Keentok, F. L. Stoddard, and F. Bekes. Basic rheology of bread dough with modified protein content and gluteninto-gliadin ratios, 744

-, F. L. Stoddard, P. W. Gras, and F. Bekes. Effects of incorporated glutenins on functional properties of wheat dough, 737

-, and --. Optimized methods for incorporating glutenin subunits into wheat dough for extension and baking studies, 731

Varavinit, S. See P. Chatakanonda, 315

Varo, P. See P. Ekholm, 562

Vasanthan, T. See J. Gaosong, 396

Vazquez-Moreno, L. See P. I. Torres, 702

Vensel, W. H. See F. M. DuPont, 607

Veraverbeke, W. S., O. R. Larroque, F. Békés, and J. A. Delcour. In vitro polymerization of wheat glutenin subunits with inorganic oxidizing agents. I. Comparison of single-step and stepwise oxidations of high molecular weight glutenin subunits, 582

, and --. In vitro polymerization of wheat glutenin subunits with inorganic oxidizing agents. II. Stepwise oxidation of low molecular weight glutenin subunits and a mixture of high and low molecular weight glutenin subunits, 589

Vergnes, B., See E. Charun, 265 Vinyard, B. T. See B. G. Lyon, 64 Virkki, L. See P. Ekholm, 562

Wahjudi, J., L. Xu, P. Wang, V. Singh, P. Buriak, K. D. Rausch, A. J. McAloo, M. E. Tumbleson, and S. R. Eckhoff. Quick fiber process: Effect of mash temperature, dry solids, and residual germ on fiber yield and purity, 640

Wahlgren, M. See U. Elofsson, 679 Wallace, A. See A. R. Wooding, 798

Wang, D., and S. R. Eckhoff. Effect of broken corn levels on water absorption and steepwater characteristics, 525

. See P. Yang, 128

Wang, F. C., D. S. Chung, P. A. Seib, and Y. S. Kim. Optimum steeping process for wet milling of sorghum, 478

Wang, P. See J. Wahjudi, 640 Wang, Y. See A. Barrett, 784

Waniska, R. D. See M. Cepeda, 489

. See E. L. Suhendro, 96

Ward, C. D. W. See W. L. Kerr, 213 Watanabe, K. See T. Adachi, 578

See H. Tang, 27

Webb, B. D. See C. J. Bergman, 454

Wieser, H. Simple determination of gluten protein types in wheat flour by turbidimetry, 48

Wilson, A. J. See A. R. Wooding, 791

Wilson, J. D. See D. B. Bechtel, 401

. See C. S. Gaines, 163

Windham, W. R. See B. G. Lyon, 64

Won, S.-Y., W. S. Choi, H. S. Lim, K.-Y. Cho, and S.-T. Lim. Viscoelasticity of cowpea starch gels, 309

Wood, D. F. See W. J. Orts, 18 See G. H. Robertson, 439

Wooding, A. R., S. Kavale, F. MacRitchie, F. L. Stoddard, and A. Wallace. Effects of nitrogen and sulfur fertilizer on the protein composition, mixing requirements, and dough strength of four wheat cultivars, 798

, A. J. Wilson, and F. L. Stoddard. Effects of nitrogen and sulfur fertilization on commercial-scale wheat quality and mixing requirements, 791

Wright, C. L. See C. K. Black, 468

Xie, X. J., and P. A. Seib. Laboratory procedure to wet-mill 100 g of grain sorghum into six fractions, 392, 696 Xu, L. See J. Wahjudi, 640

Yang, H. C. See P. A. Seib, 816

Yang, P., L. Du, D. L. Wang, B. H. Li, K. D. Rausch, P. Buriak, and S. R. Eckhoff. Effects of alkali debranning, roller mill cracking and gap setting, and alkali steeping conditions on milling yields from a dent corn hybrid,

, and S. R. Eckhoff. Reducing steep time by adding lactic acid during countercurrent steeping of corn with different initial moisture contents, 529 R. J. Shunk, A. E. Haken, Y. X. Niu, S. H. Zou, P. Buriak, S. R. Eckhoff, and M. E. Tumbleson. Yield, protein content, and viscosity of starch from wet-milled corn hybrids as influenced by environmentally induced

changes in test weight, 44 Yasui, T. See T. Sasaki, 58

. See M. Seguchi, 339

Ylinen, M. See P. Ekholm, 562 Yoshimoto, Y., J. Tashiro, T. Takenouchi, and Y. Takeda. Molecular structure and some physicochemical properties of high-amylose barley starches, 279

You, S., and S.-T. Lim. Molecular characterization of corn starch using an aqueous HPSEC-MALLS-RI system under various dissolution and analytical conditions, 303

Zheng, G. H., B. G. Rossnagel, R. T. Tyler, and R. S. Bhatty. Distribution of β-glucan in the grain of hull-less barley, 140

Zhong, Z. K., and X. S. Sun. Thermal behavior and nonfreezing water of soybean protein components, 495

Zou, S. H. See P. Yang, 44

Zweifel, C., B. Conde-Petit, and F. Escher. Thermal modifications of starch during high-temperature drying of pasta, 645

Subject Index

Page numbers of errata are in italic.

α-Amylolysis, of barley starch granules (Lauro et al), 595

α-Amylose, activity-free, relation to falling numbers (Graybosch et al), 1

Absorption, water, effect of broken corn levels on (Wang and Eckhoff), 525

Adsorption, studies of interaction between water-extractable nonstarch polysaccharides and prolamins in (Elofsson et al), 679

Aegilops spp., amylose content; starch B-granule content (Stoddard and Sarker), 445

Air classification, of barley flours (Andersson et al), 463

Alkali

-debranning corn with (Yang et al), 128

—debranning; removal of seedcoat using alkaline solution (Lazaro and Favier), 717

Amaranth, effect on serum lipids (Maier et al), 297

Amylopectin

-from high-amylose barley starch, structure (Yoshimoto et al), 279

 retrogradation kinetics of rice, enthalpy change during retrogradation (Lai et al), 272

Amylose

-Aegilops (Stoddard and Sarker), 445

-from high-amylose barley starch, structure (Yoshimoto et al), 279

Antioxidants, in rice bran; effects of commercial processing on (Lloyd et al), 551

Baking

—formula innovation to eliminate chlorine treatment of cake flour (Donelson et al), 53

—hydrolytic enzyme effect on processing and bread characteristics (Harada et al). 70

-kernel size effect on water absorption (Morgan et al), 286

Barley

—chemical composition of pearled kernels and pearling by-products (Marconi et al), 133

—extrusion cooking effect on β -glucan primary structure and water solubility (Gaosong and Vasanthan), 396

-flours; air classification of (Andersson et al), 463

—high-amylose starch, structure and physicochemical properties (Yoshimoto et al), 279

-hull-less, distribution of β-glucan in grain of (Zheng et al), 140

—lipid oxidation in relation to hydrothermal treatment (Rutgersson et al), 407 Bread

-calcium-fortified (Ranhotra et al), 293

—crumb grain characteristics; influence of additives and mixing time (Crowley et al), 370

—hydrolytic enzyme effect on processing and bread characteristics (Harada et al), 70

--phase transition, moisture redistribution and staling (Baik and Chinachoti),

—shelf-stable, textural optimization of, glyercol cotent and dough-forming technique effects (Barrett et al), 169

-vitamin E stability in (Ranhotra et al), 159

Breadmaking

-malted sorghum and wheat composite bread (Hugo et al), 428

-quality of flour affected by hot-air-drying of wheat (Tosi et al), 699

—relationship with glutenin and polymeric composition in durum wheat quality and SE-HPLC analysis in durum and bread wheat (Ammar et al), 230

Brown rice

-cakes; wheat starch effects on textural characteristics of (Orts et al), 18

-lipidemic response (Kahlon and Chow), 673

Cake, flour; baking formula innovation to eliminate chlorine treatment of (Donelson et al), 53

Calcium

—chloride; surface gelatinization with, morphological changes of starches (Koch and Jane), 115

—fortified bread; utilization as calcium carbonate or as dairy calcium (Ranhotra et al), 293

Carbodiimide, water-soluble, for cross-linking of wheat gluten (Tropini et al), 333

Cereal, adsorption studies of interaction between water-extractable nonstarch polysaccharides and prolamins in (Elofsson et al), 679

Chromatography

—effects of temperature and sonication on size distribution of proteins (Morel et al), 685

—hydrophobic interaction, in highly acetylated agarose, variability to dough rheological properties (Torres et al), 702

Cookies, semi-sweet biscuit, effect of temperature and mechanical imput during mixing on biscuit quality (Charun et al), 265

Com (see also Maize)

-alkali debranning effect (Yang et al), 128

 broken level effect on water absorption and steepwater characteristics (Wang and Eckhoff), 525

-dry-milled, solvent extraction of zein from (Shukla et al), 724

—effect of mill plate setting and number of dynamic steeping stages for IMDS process (Mehra et al), 209

—fiber and fiber oil; yield and composition affected by various acids and sulfites in steeping (Singh et al), 665

—fiber gum yields and sugar profiles; comparison of coarse and fine corn fiber (Singh et al), 560

—fiber yields and oil composition affected by growth location and hybrid variability (Singh et al), 692

—nixtamalization; dry matter loss during, processing parameter impact (Sahai et al), 255

-processing; factors affecting fiber yi eld and purity (Wahjudi et al), 640

—water absorption variability of germ and endosperm during laboratory steeping (Dailey), 721

Corn bran, bile acid binding (Kahlon and Chow), 518

Corn starch

—lipidemic response (Kahlon and Chow), 673

—molecular characterization of, using HPSEC-MALLS-RI system (You and Lim), 303

 —pellets; effect of gelatinization degree and moisture content of, on morphology and physical properties of microwave-expanded products (Lee et al), 769

—yield affected by environmentally induced test weight changes (Yang et al), 44 Cowpea, starch gels; viscoelasticity of (Won et al), 309

Debranning, corn; with alkali (Yang et al), 128

Dietary fiber

—effects of natural chelating agents on solubility of minerals in (Ekholm et al), 562

—flour, okara, affected by extrusion, enrichment in extruded products (Rinaldi et al), 237

-role in reducing serum lipids (Maier et al), 297

Dough

-bread, basic rheology of (Uthayakumaran et al), 744

-effect of temperature on tortilla properties (Cepeda et al), 489

—from three wheat flours; influence of added starch on mixing (Larsson et al), 633

-glutenin macropolymer studied by TEM (Lindsay and Skerritt), 360

-microextensibility method (Suchy et al), 39

—semi-sweet biscuit, effect of temperature and mechanical imput during mixing (Charun et al), 265

—wheat, functional properties of, affected by incorporated glutenins (Uthaya-kumaran et al), 737

—wheat, optimized methods for incorporating glutenin subunits into (Uthayakumaran et al), 731

-wheat flour, effects of laccase and ferulic acid on (Labat et al), 823

Emulsion, water, oil in, stabilized with peptides (Linarès et al), 414

Endosperm

-proteins in wild and opaque-2 maizes (Landry et al), 620

 water absorption variability of, during laboratory steeping of yellow corn hybrid (Dailey), 721

Environment, effect on oat milling characteristics and groat hardness (Doehlert and McMullen), 148

Enzymes, hydrolytic, effect on processing and bread characteristics (Harada et al), 70

Erratum, vol. 77, no. 3, 39, 496

Ethanol, process of producing, from corn (Wahjudi et al), 640

Extrudates, intermediate-moisture, effects of formulation on textural stability of (Barrett et al), 784

Extrusion

- —of cross-linked hydroxypropylated corn starches; morphological and molecular characterization (McPherson and Jane), 326
- —of cross-linked hydroxypropylated corn starches; pasting properties (McPherson et al), 320
- —temperature and screw speed effect on oil absorption characteristics of a multigrain extrudate during frying (Osman et al), 101
- —of wheat flour, okara, effects on dietary fiber and isoflavone contents (Rinaldi et al), 237
- —of wheat flour; changes in disulfide and sulfhydryl contents (Anderson and Ng), 354

Extrusion cooking

- —effect on β-glucan primary structure and water solubility, from regular and waxy barley (Gaosong and Vasanthan), 396
- —pet food twin-screw, residence time distribution and barrel fill (Mahungu et al), 220

Falling number, determination of oats (Kaarlehto and Salovaara), 177

- —com, coarse and fine comparison, for corn fiber gum yields and sugar profiles (Singh et al), 560
- -recovery prior to ethanol production (Wahjudi et al), 640
- Flavor, term used to describe combined results of taste, odor, and textural feelings (Sayaslan et al), 249

Flour

- —Asian noodles from hard white and hard red wheat flours, comparison (Seib et al), 816
- -barley, air classification of (Andersson et al), 463
- -breadmaking quality affected by hot-air-dryi ng of wheat (Tosi et al), 699
- -cake, baking formula innovation to eliminate chlorine treatment of (Donelson
- —composition; genotypic and environmental effects (Mikhaylenko et al), 507
- —cowpea, effect on functionality and physicochemical properties of (Kerr et al), 213
- -effects of extrusion conditions on flour proteins (Anderson and Ng), 354
- —enriched with okara dietary fiber and isoflavones in extruded products (Rinaldi et al), 237
- -kernel size effect on water absorption (Morgan et al), 286
- -protein; size distribution by SE-HPLC (Morel et al), 685
- -rice, modeling starch gelatinization kinetics of (Ojeda et al), 145
- -turbidity for simple determination of gluten protein types in (Wieser), 48
- —polylactic acid starch-based loose-fill packaging, functional properties of (Fang and Hanna), 779
- -stabilized by peptides derived from gluten (Linarès et al), 414

Fractionation, of flour protein (DuPont et al), 607

Gas chromatography, method for 2-acetyl-1-pyrroline and hexanal in milled rice (Bergman et al), 454

Gelatinization

- -of cross-linked rice starch (Chatakanonda et al), 315
- -effect of amylose content on (Sasaki et al), 58
- Glass transition, DSC and DMA analyses for compatibility of rice starch and pectins (Lai et al), 544

Gliadin

- -enzymatic polymerization (Larré et al), 121

β-Glucan

- -in barley pearling by-products (Marconi et al), 133
- -distribution in grain of hull-less barley (Zheng et al), 140
- —from regular and waxy barley; extrusion cooking effect on primary structure and water solubility (Gaosong and Vasanthan), 396

Gluten

- —hydrolysates; effect of soluble and insoluble fractions on emulsion and foam stability (Linarès et al), 414
- —polymeric composition in durum and bread wheat, relationship in breadmaking (Ammar et al), 230
- —proteins; accessibility of amino groups studied by combination of chemical labeling and immunochemical detection (Scardone et al), 602
- -proteins; contribution to dough rheological properties (Torres et al), 702
- -protein types determined by turbidity, in wheat flour (Wieser), 48
- ---separation; effect of gluten structure on displacement of starch (Robertson et al), 439
- -structure (Lindsay and Skerritt), 360
- -transglutaminase-modified, rheological properties of (Larré et al), 121

Glutenin

- —allelic composition; breadmaking quality of durum wheat (Ammar et al), 230
- -changes in solubility, size of polymers (Larré et al), 121
- incorporated, effects on functional properties of wheat dough (Uthayakumaran et al), 737
- —in vitro polymerization of HMW-GS and LMW-GS isolated from wheat (Veraverbeke et al), 582, 589
- —rapid analysis of molecular size distribution by SE-HPLC (Larroque and Bekes), 451
- —subunits; methods for incorporating into wheat dough (Uthayakumaran et al), 731
- Grain sorghum, lab procedure to wet-mill into six fractions (Xie and Seib), 392. 696
- Grits, production; scarification and degermination of sorghum for (Suroso et
- Gum, corn fiber, comparison of coarse and fine corn fiber (Singh et al), 560

Hardness

- —associations of starch gel hardness, granule size, waxy allelic expression, thermal pasting, milling quality, and kernel texture of 12 soft wheat cultivars (Gaines et al), 163
- -groat, genotypic and environmental effects on (Doehlert and McMullen), 148
- parameters and composition, correlation, Argentine maize races (Robutti et al), 24
- —pentosan fraction and end-use quality of wheat relationship (Bettge and Morris), 241

High molecular weight

- -glutenin subunits; isolated from durum wheat (Capocchi et al), 105
- —glutenin subunits; temperature-induced changes in rheology and size distribution of gluten proteins from wheat near-isogenic lines (Lefebvre et al), 193
- -glutenins; atomic force microscopy images (Humphris et al), 107

HPLC

- -analysis of isoflavone contents in flour and okara (Rinaldi et al), 237
- —size exclusion analysis of semolina and whole meal extracts (Larroque et al), 448
- —size-exclusion rapid analysis of molecular size distribution for wheat proteins (Larroque and Bekes), 451
- HPSEC, to determine molecular characterization of corn starch (You and Lim), 303
- Hydrophobicity, surface index relation of soluble gluten proteins to dough rheological properties (Torres et al), 702
- Hydrothermal treatment, influence on lipid oxidation (Rutgersson et al), 407

Image analysis

- —for appearance assessment of oriental noodles as function of flour refinement and noodle type (Hatcher and Symons), 181
- —of Asian noodle appearance; impact of hexaploid wheat with red seed coat (Hatcher and Symons), 388
- —for assessing sprout damage influence on Asian noodles (Hatcher and Symons),

-of isolated wheat starch (Bechtel and Wilson), 401

Immunochemistry, for detection of chemical probe used to label primary amino groups in gluten proteins (Scardone et al), 602

Instructions to authors, iv

Instruments and instrumentation

- -determining kernel vitreousness using NIR (Dowell), 155
- -torque measurement during milling with a micromill (Pujol et al), 421

Lactic acid, addition during corn steeping to reduce steep time (Yang and Eckhoff), 525

Leavening, amount and type of acids in tortilla properties (Cepeda et al), 489 Lipidemic response, of hamsters to rice bran, white and brown rice, processed corn starch (Kahlon and Chow), 673

Lipids

- —degradation in stored rice; measurement with TLC/FID sysem (Nishiba et al), 223
- —effect of addition on α -amylolysis of barley starch granules (Lauro et al), 595
- -free, NIR transmittance estimation of (Ohm and Chung), 556
- —oxidation; in barley during cereal processing and storage (Rutgersson et al), 407

Maize (see also Corn)

- —endosperm; improved method for sequential extraction of proteins (Landry et al), 620
- —grouping and identification of races by principal component analysis of RP-HPLC zein data (Robutti et al), 91
- -quality factors, in Argentine races (Robutti et al), 24
- —quality protein, nutritional value of weaning foods (Barragán Delgato and Serna Saldivar), 652
- —waxy starch-sugar extrudates, sugar effect on retrogradation (Farhat et al), 202

Malt, sorghum, use in breadmaking (Hugo et al), 428

Methods

- -for 2-acetyl-1-pyrroline and hexanal in milled rice (Bergman et al), 454
- -digital image analysis of isolated wheat starch (Bechtel and Wilson), 401
- -extensibility, small scale (Suchy et al), 39
- —to group and identify maize races by principal component analysis of zein RP-HPLC data (Robutti et al), 91
- -noodle color affected by variables (Morris et al), 77

Microencapsulated high-fat powders, effect on dough rheology (O'Brien et al), 111

Microwave, expanded products; effect of gelatinization degree and moisture content of, on morphology and physical properties of (Lee et al), 769 Millet, alkali debranning of (Lazaro and Favier), 717

Milling

- —developing agreement between very short flow and longer flow test wheat mills (Gaines et al), 187
- —effect on functionality and physicochemical properties of cowpea flour (Kerr et al), 213
- -energy consumption and particle size reduction (Pujol et al), 421
- -kernel size effect on flour water absorption (Morgan et al), 286
- —of oats; genotypic and environmental effects on (Doehlert and McMullen), 148

Mineral, elements in oat bran and oat flakes; effects of chelating agents on solubility (Ekholm et al), 562

Modeling, of water sorption kinetics in spaghetti (Del Nobile and Massera), 615 Moisture, influence on rough rice respiration rate (Dillhunty et al), 541

Mung bean, starch; effect of annealing on gel property (Chung et al), 567

Near-infrared transmittance spectroscopy, for classification of single- and doublemutant corn endosperm genotypes (Campbell et al), 774

NIR, of individual wheats; protein content of bulk wheat (Delwiche and Hruschka), 86

NIR spectroscopy

- -classification of durum vitreousness (Dowell), 155
- —of free lipid content and its glycolipid and digalactosyldiglyceride contents (Ohm and Chung), 556
- -rice quality by (Barton et al), 669

Nitrogen

-effects of on commercial-scale wheat quality and mixing requirements

(Wooding et al), 791

—fertilizer effect on protein composition, mixing, and dough strength of four wheat cultivars (Wooding et al), 798

Noodles

- —Asian, from hard white and hard red wheat flours, comparison (Seib et al), 816
- —Asian, image analysis for assessing sprout damage influence on (Hatcher and Symons), 380
- —Asian, image analysis of, impact of hexaploid wheat with red seed coat (Hatcher and Symons), 388
- --from food sorghum; cooking characteristics and quality (Suhendro et al),
- oriental, appearance assessment as function of flour refinement and noodle type by image analysis (Hatcher and Symons), 181
- -wheat of white salted, quality (Black et al), 468
- -yellow alkaline, effect of variables (Morris et al), 77

Oat

- —falling number; determination of, effect of kiln drying on (Kaarlehto and Salovaara), 177
- genotypic and environmental effects on milling characteristics (Doehlert and McMullen), 148
- --proteinases of germinated (Mikola and Jones), 572

Oat bran

- -bile acid binding (Kahlon and Chow), 518
- -effect on serum lipids (Maier et al), 297
- Oil, absorption characteristics of a multigrain extrudate during frying, temperature and screw speed effect (Osman et al), 101
- Oxidation, of HMW-GS and LMW-GS with inorganic oxidants (Veraverbeke et al), 582, 589

Pasta

- -cooked, starch retrogradation in (Riva et al), 433
- —drying; thermal modifications of starch during high-temperature drying (Zweifel et al), 645
- -products; peroxidase activity effect on (Fraignier et al), 11
- -unconventional ingredients; barley pearling by-products (Marconi et al), 133
- Pearl millet, decortication; nutritional value of weaning foods (Barragán Delgato and Serna Saldivar), 652
- Pentosans, relation to grain hardness and end-use quality of wheat (Bettge and Morris), 241
- Peroxidase, activity and localization in durum wheat (Fraignier et al), 11
- Physicochemical properties, of starches in fractions of waxy barley grain (Tang et al), 27
- Polysaccharides, water-extractable nonstarch, interaction between prolamins in cereals (Elofsson et al), 679
- Processing, commercial, effects on antioxidants in rice bran (Lloyd et al), 551
- Prolamins, interaction between water-extractable nonstarch polysaccharides in cereals (Elofsson et al), 679
- Protease, interference in analysis of sorghum proteins (Huang et al), 343
 Protein
- —amino acid composition in sorghum, maize, and pearl millet (Barragán Delgato and Serna Saldivar), 652
- -bulk wheat content, from NIR (Delwiche and Hruschka), 86
- -changes due to extrusion processing (Anderson and Ng), 354
- -effect on durum NIR spectra (Dowell), 155
- -fraction in maize endosperm (Landry et al), 620
- —ω-gliadin sequence and characterization (DuPont et al), 607
- -instability of SE-HPLC extracts (Larroque et al), 448
- -protease interference in analysis of sorghum proteins (Huang et al), 343
- -quality; micro SDS sedimentation test (Mikhaylenko et al), 507
- -quality; SDS sedimentation volume (Black et al), 468
- -rapid SE-HPLC analysis (Larroque and Bekes), 451
- -SE-HPLC durum and bread wheat (Ammar et al), 230
- -size distribution and extractability by SE-HPLC (Morel et al), 685
- -thiamin-binding from wheat germ, characterization of (Adachi et al), 578

Proteinases, of germinated oats that hydrolyze oat globulins (Mikola and Jones), 572

Rapid Visco Analyser

-effect of temperature profile on viscosity (Batey and Curtin), 754

-mixing system; average shear rates in (Lai et al), 714

Retrogradation

-kinetics; effect of molecular property on rice amylopectin (Lai et al), 272

-mechanism of rice starch (Tako and Hizukuri), 473

-starch, in cooked pasta and rice (Riva et al), 433

Reviewers, acknowledgment of, iii

Rheological behavior, small and large deformation (Larsson et al), 633 Rheological properties

-of gluten modified by transglutaminase (Larré et al), 121

-sorghum malt dried under various conditions (Hugo et al), 428

Rheology

-basic, of bread dough (Uthayakumaran et al), 744

-dough, effects of dairy ingredients and added fat on (O'Brien et al), 111

-extensigraph and TA.XT2, comparison of (Suchy et al), 39

—semi-sweet biscuit dough, effect of temperature and mechanical imput during mixing (Charun et al), 265

Rice

—chalky kernels; physicochemical characteristics and effects on sensory quality of cooked rice (Kim et al), 376

—chalky and translucent grains; differences in starch composition and structure and cooking properties (Lisle et al), 627

—cooked long-grain, sensory quality of, factors affecting (Meullenet et al), 258

-cooked, starch retrogradation in (Riva et al), 433

-cooked, texture evaluation (Lyon et al), 64

-drying; development of brown rice state diagram (Perdon et al), 708

—effect of blast and sheath blight on kernel physical properties and head rice yield (Candole et al), 535

-flour; modeling starch gelatinization kinet ics of (Ojeda et al), 145

 instrumental assessment of cooked texture characteristics, method for breeders (Meullenet et al), 512

-method for 2-acetyl-1-pyrroline and hexanal (Bergman et al), 454

—prediction of cooked using extrusion and compression tests in conjunction with spectral stress strain analysis (Sitakalin and Meullenet), 501

-quality by spectroscopic analysis (Barton et al), 669

-respiration rate; effect of moisture content and temperature (Dillhunty et al), 541

-simple measurement of lipid degradation during storage (Nishiba et al), 223

-starch; compatibility of by glass transition (Lai et al), 544

-starch; gelatinization (Tako and Hizukuri), 473

—starch; gelatinization and retrogradation of cross-linked (Chatakanonda et al), 315

Rice bran

-antioxidants; effects of commercial processing on (Lloyd et al), 551

-bile acid binding (Kahlon and Chow), 518

-lipidemic response (Kahlon and Chow), 673

RP-HPLC, grouping and identification of maize races by principal component analysis of zein data (Robutti et al), 91

SDS-PAGE, of flour proteins from various extrusion conditions (Anderson and Ng), 354

SE-HPLC, comparison between durum and bread wheat, relationship with breadmaking quality (Ammar et al), 230

Semolina, size exclusion analysis (Larroque et al), 448

Sorghun

-alkali debranning of (Lazaro and Favier), 717

—malted, nutritional value of weaning foods (Barragán Delgato and Serna Saldivar), 652

-malt used in breadmaking (Hugo et al), 428

-noodles; cooking characteristics and quality of (Suhendro et al), 97

-scarification and degermination of, for grits production (Suroso et al), 808

—wet milling; starch yield and color affected by steeping parameters, temperature, time, and concentration of media (Wang et al), 478

Soy, protein; effect of moisture content and sample preparation method on thermal properties (Zhong and Sun), 495

Spaghetti, modeling of water sorption kinetics in, during overcooking (Del Nobile and Massera), 615

Starch

--α-amylolysis of (Lauro et al), 595

—associations of starch gel hardness, granule size, waxy allelic expression, thermal pasting, milling quality, and kernel texture of 12 soft wheat cultivars (Gaines et al), 163

--biopolymer, made of mainly amylose and amylopectin (Sayaslan et al), 249

-B-type granules, Aegilops (Stoddard and Sarker), 445

 cationized waxy maize granules; enzymatic and acidic hydrolysis (Manelius et al), 345

-com, single- and double-mutant genotype classification (Campbell et al), 774

 cross-linked hydroxypropylated, morphological and molecular characterization of extruded (McPherson and Jane), 326

—cross-linked hydroxypropylated, pasting properties of extruded (McPherson et al), 320

-effect on durum NIR spectra (Dowell), 155

—foam extruded with normal and waxy corn starches blended with polylactic acid polymer (Fang and Hanna), 779

-gelatinization kinetics; modeling of, in milled rice flour (Ojeda et al), 145

-gels; viscoelasticity of (Won et al), 309

—granules in rice; compound amyloplasts containing a number of (Lisle et al), 627

—high-amylose barley, structure and physicochemical properties (Yoshimoto et al). 279

-influence on mixing of wheat flour dough (Larsson et al), 633

-internal structure of (Seguchi et al), 339

—morphological changes of, by surface gelatinization with calcium chloride (Koch and Jane), 115

-mung bean, effect of annealing on gel property (Chung et al), 567

—oat, effect of kiln drying on pasting properties (Kaarlehto and Salovaara), 177

-pasting properties; amylose content effect (Sasaki et al), 58

-and poly(lactic acid) composites; physical p roperties of (Ke and Sun), 761

-retrogradation in cooked pasta and rice (Riva et al), 433

—role of amylopectin structure in, retrogradation kinetics in various starchcontaining systems (Lai et al), 272

—size distribution determined by digital image analysis (Bechtel and Wilson), 401

-starch granule size distribution (Black et al), 468

—thermal modifications of, during high-temperature drying of pasta (Zweifel et al), 645

et al). 44

-waxy granules, susceptibility to mechanical damage (Bettge et al), 750

—waxy maize, structure of, α-amylolysis pattern, intermediate pattern (Bertoft et al), 657

---wheat, effects on textural characteristics of puffed brown rice cakes (Orts et al), 18

State diagram, of rice; glassy state transition and rice drying on development of (Perdon et al), 708

Steeping

—acid and sulfite effect on yields and composition of corn fiber and corn fiber oil (Singh et al), 665

-of corn; reducing steep time by adding lactic ac id (Yang and Eckhoff), 525

-effect of broken corn levels on (Wang and Eckhoff), 525

—sorghum; starch yield and color affected by steeping parameters, temperature, time, and concentration of media (Wang et al), 478

—of yellow corn hybrid; variability of water absorption of germ and endosperm during (Dailey), 721

Storage, temperature and duration affecting sensory quality of cooked longgrain rice (Meullenet et al), 258

Storage protein, characterization of thiamin-binding protein from wheat germ (Adachi et al), 578

Sugar, effect on retrogradation of waxy maize starch-sugar extrudates (Farhat

Surface gelatinization, of starch granules beginning from the periphery and inward (Koch and Jane), 115

Temperature

-influence on rough rice respiration rate (Dillhunty et al), 541

—mixing, effect on semi-sweet biscuit dough characteristic and biscuit quality (Charun et al), 265

Textur

—of cooked rice: instrumental assessment, method for breeders (Meullenet et al), 512

- prediction using extrusion and compression tests in conjunction with spectral stress strain analysis (Sitakalin and Meullenet), 501
- -profile of cooked rice (Lyon et al), 64
- Thermal properties, denaturationn temperature and enthalpy obtained from differential scanning calorimetry (Zhong and Sun), 495
- Tortilla, effect of leavening acids and dough temperature (Cepeda et al), 489

Viscoelasticity, of cowpea starch gels (Won et al), 309

Viscosity

- —effect of varying conditions in Rapid Visco Analyser (Batey and Curtin), 754
- -gel, effect of compatibility on (Lai et al), 544
- -liquefied weaning foods (Barragán Delgato and Serna Saldivar), 652
- -pasting property by RVA (McPherson et al), 320
- Vitamin E, added to bread (Ranhotra et al), 159

Water, nonfreezing, bound water of soy protein (Zhong and Sun), 495

Wet-milling, sorghum; starch yield and color affected by steeping parameters, temperature, time, and concentration of media (Wang et al), 478

Wheat

- —associations of starch gel hardness, granule size, waxy allelic expression, thermal pasting, milling quality, and kernel texture of 12 soft wheat cultivars (Gaines et al), 163
- —developing agreement between very short flow and longer flow test wheat mills (Gaines et al), 187
- —durum, breadmaking quality, gluten polymeric composition (Ammar et al),
- -durum, peroxidase activity (Fraignier et al), 11
- —durum, Single-Kernel Characterization System for testing and quality prediction (Sissons et al), 4
- —durum, wheat high molecular weight glutenin subunits isolated from (Capocchi et al), 105
- —end-use quality, grain hardness, and pentosan fraction relationship (Bettge and Morris), 241
- —flour; appearance assessment of oriental noodles as function of flour refinement and noodle type by image analysis (Hatcher and Symons), 181
- —flour, hydrolytic enyme effect on processing and bread characteristics (Harada et al), 70
- —flour, turbidity for simple determination of gluten protein types in (Wieser), 48
- -flour doughs; effects of laccase and ferulic acid on (Labat et al), 823

- —flour lipid extracts for NIR transmittance estimation of free lipid content and its glycolipid and digalactosyldiglyceride contents (Ohm and Chung), 556
- —germ; characterization of thiamin-binding protein from (Adachi et al), 578 —gluten; cross-linking using water-soluble carbodiimide (Tropini et al), 333
- —gluten and starch; factors affecting separation (Robertson et al), 439
- —hardness; mechanical resistance to grinding with a micromill (Pujol et al), 421
- —hexaploid with red seed coat; impact on image analysis of Asian noodles (Hatcher and Symons), 388
- -hot-air-drying, effect on breadmaking quality of flour (Tosi et al), 699
- -kernel size effect on flour water absorption (Morgan et al), 286
- -landrace; white salted noodle quality characteristics in (Black et al), 468
- —near-isogenic lines; glutens from, temperature-induced changes in rheology and size distribution (Lefebvre et al), 193
- —nitrogen and sulfur fertilizer effect on protein composition, mixing, and dough strength of (Wooding et al), 798
- -noodles, effect of variables (Morris et al), 77
- -protein content of bulk, NIR of individual wheats (Delwiche and Hruschka), 86
- —quality and mixing requirements; affected by nitrogen and sulfur fertilization (Wooding et al), 791
- —quantification of durum vitreousness using NIR spectroscopy (Dowell), 155
- —size distribution of starch granules determined by digital image analysis (Bechtel and Wilson), 401
- —spring, environmental effects on experimental baked product quality (Mikhaylenko et al), 507
- -starch; effects on textural characteristics of puffed brown rice cakes (Orts et al), 18
- -starch; internal structure studied by KI/I, solution (Seguchi et al), 339
- -starch pasting properties; amylose content effect (Sasaki et al), 58
- -waxy, falling numbers of (Graybosch et al), 1
- Wheat bran, bile acid binding (Kahlon and Chow), 518

Zein

- —biodegradable sheets plasticized with oleic acid; thermal behavior (Budi Santosa and Padua), 459
- —from dry-milled corn; solvent extraction of (Shukla et al), 724
- —grouping and identification of maize races by principal component analysis of RP-HPLC data (Robutti et al), 91

